

# The Effects of Pollution Reduction on a Wild Trout Stream

## Final Report: 2005 - 2008



Spring Run

Dumpling Run



**April 2009**

Prepared by W. Neil Gillies, Cacapon Institute

## TABLE OF CONTENTS

<b>Introduction</b> .....	3
<b>Partners</b> .....	5
<b>Methods</b> .....	6
<b>Hatchery Upgrade</b> .....	7
<b>Results</b> .....	9
Flow .....	9
Water quality .....	9
How did water quality vary over time? .....	12
How did loads of key parameters vary over time? .....	14
<b>Benthic Macroinvertebrate &amp; Periphyton Analysis</b> .....	16
<b>Fisherman Survey</b> .....	18
<b>DISCUSSION</b> .....	20
<b>Erosion Control Projects</b> .....	21
<b>Outreach</b> .....	22
<b>Priority Watershed Planning</b> .....	22
<b>Year Four expectations</b> .....	22
<b>Appendix 1. Water Quality Statistics by Year</b> .....	23
Appendix 2. Laboratory Methods for Water Quality Parameters. ....	29
Appendix 3. WV Save Our Streams Macroinvertebrate Assessment July 2003 .....	30
<b>Appendix 4. Assessing the Condition of the Macroinvertebrate Communities of Spring Run</b> ..	31

## **The Effects of Pollution Reduction on a Wild Trout Stream Final Report: 2008**

### **Introduction**

Spring Run is a unique aquatic resource in the Potomac Highlands region of West Virginia. Unlike many small headwater streams that tend to go dry, it is fed by the largest spring in the region, with discharge typically ranging from 3000-3500 gallons per minute. With a temperature of ~53 °F at the spring and a pH of ~8, aquatic conditions are ideal for trout and the aquatic insects they eat. Spring Run flows about two miles from the spring source to its confluence with South Mill Creek, which is about four miles from the South Branch of the Potomac River. Spring Run has no tributaries. Much of the stream is shallow, and does not provide the complex habitat that trout need - but that is not the case in a one mile section in the middle of the Run.

Since the early 1960's, landowner's have issued permits for fly fishing, catch-and-release on about one mile of Spring Run. Landowners and other interested parties have installed and maintained various structures to form pools and overhead cover that provide hiding and feeding habitat for trout. Spring Run is recognized as one of the best "wild" rainbow trout fisheries in West Virginia. Friends of Springs Run's Wild Trout, was formed in 1996 to restore structure to Spring Run following flooding in 1996.

In the last few years, however, fishermen have noted a decline in the fishery. Emergence of the mayfly, Ephemerellidae (sulfurs) largely disappeared in the late 1990s. The number of large trout (14" and above) has decreased and trout in the 11-13" range have also declined in abundance. The population of trout is considerably lower in the lower reach of the three-fourths mile section. Algae formation is heavy in the upper reach of the catch-and-release section, much heavier than in the past, and algae reforms soon after washout by high water.

Spring Run is rich in nutrients, delivered largely in effluent from the Spring Run Trout Hatchery (SRH) which is located about one-third mile upstream from the upper end of the fly fishing section and about one-fourth mile below the spring. (SRH is a rearing facility; trout are not spawned there). In recent years, however, SRH has been producing more rainbow and "golden trout" for stocking West Virginia streams, and it seems that the effluent stream now may be a problem for the health of Spring Run. WVDEP issued a citation for violation of the Spring Run Trout Hatchery NPDES permit in January 2004, specifically for discharging excess biochemical oxygen demand (BOD) and total suspended solids (TSS). WVDNR, which operates SRH, installed an effluent treatment system at the facility to meet their permit requirements. It became operational in June 2007.

The prospective installation of effluent treatment at SRH provided a unique opportunity to address a number of issues of both regional and national significance:

1. Will the hatchery effluent treatment process significantly reduce nutrient discharge? Fish hatcheries throughout the country produce nutrient-rich effluents of concern to receiving waters. This study will evaluate the downstream result of effluent reduction of BOD and TSS, as well as nutrients, from a small but high throughput point source. The results of renovation at SRH and this study will provide important information to the WV Potomac Tributary Strategy point source innovation process.
2. What are the biological impacts of Spring Run's high nutrient levels, and how is the biota affected by reductions in nutrients, TSS and BOD following hatchery upgrades? This issue is of importance to the nutrient criteria development process that WV and the other 49 states are currently struggling through, as one of the key questions is: "what does nutrient impairment look like?"

3. Is the wild trout population in Spring Run being harmed by hatchery effluent, and does improvement in that effluent improve the trout fishery?
4. Is the benthic invertebrate population in Spring Run being harmed by hatchery effluent, and does improvement in that effluent improve diversity? Spring Run fishermen have noted the loss in recent years of a certain family of mayflies, the Ephemerellidae (Spiny crawler mayfly) that used to emerge regularly in the springtime. Also, WV DEP's Tim Craddock completed a benthic assessment of Spring Run in 2002, and found the lower part of the fly fishing section to be dominated by Chironomidae (midge) larvae, a group often indicative of pollution by organic waste.
5. Why do trout, especially larger fish, favor the upper part of the fly-fishing section? Why has the density-center of the trout population moved upstream in recent years? Is there a relationship between distribution of benthic invertebrates in the stream and trout distribution? If the Ephemerellidae mayflies and other pollution sensitive macroinvertebrates rebound after the hatchery effluent is treated, will the trout population improve also? In particular, are trout avoiding areas they used to frequent that are now dominated by midge larvae? If upgrades to the hatchery reduce organics in the stream and also the midge populations, will trout return to those areas? If that turns out to be true, and we could demonstrate that it is true, that would buttress public acceptance of benthic invertebrate stream assessments.

Overall, this project has the potential to address many questions beyond the five questions identified above.

This progress report will provide an overview of two years of baseline data and the nearly two years of post-treatment data. Additional sampling of benthic macroinvertebrates and the fish community are planned, but unfunded at this time.

### Timeline

The following figure provides an approximate timeline for significant events related to the fishery in Spring Run.

<b>SPRING RUN TIMELINE (APPX.)</b>																
PAST TO 1930	1930	1940	1950	1960 - 1989	1990- 1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
BROOK TROUT FISHERY																
	RAINBOWS INTRODUCED															
	RAINBOW TROUT FISHERY															
	SPRING RUN HATCHERY (ON-LINE 1952)															
			SPRING RUN STOCKED TILL MID 1980's													
			SPRING RUN PRIVATELY MANAGED C&R													
					NOTICE "SULFUR" HATCH AND "YELLOW SALLIES" MISSING											
								RECOGNIZE DECLINE IN RAINBOW TROUT FISHERY								
								HATCHERY CITED FOR PERMIT VIOLATIONS								
												SPRING RUN ASSESSMENT PROJECT				
NOTES: TIMELINE APPROXIMATE.												STOP RELEASE OF TROUT FROM VIEWING POND				
EARLY DATES BASED ON ANECDOTAL ACCOUNTS FROM A FEW PEOPLE.												NO TROUT ESCAPEES FROM HATCHERY				
SPRING RUN STOCKING OCCURRED FROM HATCHERY OPENING THROUGH MID 1980s												TREATMENT SYSTEM ON-LINE JUNE 07				

## Partners


Friends of Spring Run's Wild Trout, Cacapon Institute (CI), the WV Conservation Agency (WVCA), WV Department of Agriculture (WVDA), WV Division of Natural Resources (WVDNR), WV Department of Environmental Protection (WVDEP), and the Freshwater Institute are partnering in this study. This project is funded primarily by West Virginia Conservation Agency's participation through the Chesapeake Bay Program. An associated erosion reduction project is funded through a Friends of Spring Run's Wild Trout 2005 Stream Partners Grant. Additionally, a home school group is monitoring the lower portion of Spring Run on a regular basis.

WVDA, WVDEP and WVDNR are all contributing in-kind services to the project. WVDA is collecting water samples, taking flow measurements, and performing field and laboratory water quality analyses. WVDEP is participating in collections of benthic invertebrate and periphyton and helping to cover the costs of analysis. WVDNR is performing fish surveys and Friends of Spring Run's Wild Trout is providing information on size and location of trout caught and released by permitted fly fisherman, and supplied equipment and labor for erosion-sediment control work..

The Freshwater Institute provided guidance to WVDNR on treatment methods for their effluent and is providing technical guidance for the project. WVCA is acting as project coordinator. Cacapon Institute has overall technical oversight for the project, will participate in field work, and will, in cooperation with partnering organizations, be responsible for data analysis and production of annual reports.

## Methods

The project has two experimental components, an upstream/downstream design in Spring Run, and a control/experimental design that includes Dumpling Run, another spring fed stream nearby. Both streams have their origins in the same geology: limestone (Helderberg and Tonoloway/Wills Creek) and sandstone (Oriskany, McKenzie) formations. Spring Run flows off the ridge to the northwest into South Mill Creek, a tributary of the South Branch of the Potomac River. Dumpling Run flows east into the South Fork of the South Branch of the Potomac River.



The upstream/downstream part includes three sites in Spring Run: the first site is near the spring upstream of the hatchery; the second site is near the upper end of the fly fishing stream section; and the third is near the lower end of the fly fishing section. There are two sites on Dumpling Run, one just below the spring, the other some distance downstream. Overall, this design allows within stream and between stream comparisons. Under most conditions of flow the springs constitute the main source of water in both streams, but both streams also have periodic surface flow entering the main channel upstream of the spring. Due to unanticipated delays in construction of the effluent treatment system, the baseline period of data collection lasted for approximately two years (April 2005-May, 2007).

Water chemistries were collected monthly from April through September, typically on Wednesday. We chose to avoid collections on Mondays at the time of the pre-treatment hatchery cleanout because the "biosolids from the aquaculture effluent are notoriously patchy and difficult to characterize in sampling. . . . my thoughts on the nutrients is to focus on the residual chronic impacts, not the pulse of the cleaning plume"


(Joe Hankins, Freshwater Institute, personal communication). However, due to scheduling requirements, samples in September 2006 were collected on a Monday during the cleanout.

Water quality parameters include nitrogen in the forms of ammonia-nitrogen, nitrate/nitrite, total Kjeldahl nitrogen, total nitrogen (the sum of nitrate/nitrite and TKN), soluble reactive phosphorus, total phosphorus, total suspended solids (TSS), biochemical oxygen demand (BOD<sub>5</sub>), and basic field parameters (pH, temperature, conductivity) (see Appendix 2 for laboratory methods). Flow measurements are collected at the same time as water samples at one site in each stream. This work is done primarily by the WVDA.

Benthic invertebrate and periphyton samples are collected twice each year at all sites, in spring and autumn, according to the standard protocols in use by the WVDEP. WVDEP format Rapid Bioassessment Protocol habitat analyses is conducted once each year. WVDEP and Cacapon Institute are primarily responsible for this fieldwork.

WVDNR will conduct electro shocking fishery assessments, and the permitted fly fishermen of Spring Run have been enlisted to record information on size and location of trout caught and released.

The methods used to analyze water quality data were graphical and statistical. Data distributions were displayed using box plots (sample box-plot at right), which are useful for side-by-side visual comparisons of data distributions. One way analysis of variance (ANOVA) was run on rank transformed data for comparison of median concentration distributions. An alpha value of 0.05 was used as the threshold for statistical significance. If a significant difference among group medians was detected, Tukey's multiple comparison test was used on the rank transformed data to determine where differences were located (Helsel and Hirsh, 1992). Statistics were calculated using JMP Statistical Discovery Software (version 4.0.2). Summary statistics and raw data are provided in Appendix XX.





### Hatchery Upgrade (per Mike Shingleton, WVDNR)

The Spring Run Trout Hatchery effluent treatment system became fully operational on June 4, 2007. An automated composite sampler was installed as part of the hatchery renovation and was first used in April.

Cleaning process: Blocking weirs are placed in front of the quiescent zone of each raceway prior to cleaning; a standpipe is removed and the quiescent zone is brushed cleaned; 3-4 raceways are cleaned at a time; the wastewater from the quiescent zones is piped into the clarifier while water from raceways not being cleaned is discharged to Spring Run. The clarifier is filled to its holding capacity but is not allowed to overflow. Wastewater in the clarifier is allowed to settle 24-48 hours and the clarified water is then decanted and mixed with hatchery water back into Spring Run. The sludge remaining in the clarifier is pumped to the sludge holding tank for later disposal by land application. The decanting process of discharging water from the clarifier occurs 2-3 times per week and lasts from 1 1/2 to 2 hours each time. The following figures show reductions in total nitrogen, total phosphorus, BOD and TSS due to the treatment process. These results are based on composited samples collected during the cleanout process. Prior to the upgrade, these samples were collected during the actual cleanout. After the treatment system became operational, samples were collected during decanting.

Results of composite sample analysis of Spring Run Hatchery effluent during cleanout before and after treatment system became operational. The "before" period was from September 2006 through May 2007. The "after" period was from June through September 2007.






## Results

The effluent treatment system at the hatchery became operational in June 2007, leaving four months (June through September) of post-operational data in 2007 and a full six months in 2008. Appendix 1 provides water quality statistics by year. Appendix 4 will provide detailed benthic invertebrate statistics once data are available.

**Flow.** Stream flows on sampling days for the most downstream sites in Spring Run and Dumpling Run are given in Figure 1. Flow in Spring Run was always at least twice as high as in Dumpling Run. Sampling day flows in 2005 were much more variable than in following years, and the streams were notably low between July and September in 2007, and between July and September 2008 – six of the ten months of post-treatment data. Due to the relationship between flow and the water quality parameters measured, these extended low flow conditions have an impact on the ability to draw statistical conclusions as to effects of the treatment system.



**Water Quality.** Median values for pH, conductivity and dissolved oxygen by site and by year are provided in Table 1. These data indicate the source water in each stream was very similar, and these median values varied narrowly across all sites and all years. Both streams are alkaline, with moderately high conductivity, and high dissolved oxygen levels. No difference between pre-treatment (2005 and 2006) and post-treatment (2007) periods was evident.

**Table 1. Median pH, conductivity, and dissolved oxygen by site and year.**

Site	Median pH				Median Conductivity				Median Dissolved Oxygen (mg/L)			
	2005	2006	2007	2008	2005	2006	2007	2008	2005	2006	2007	2008
Dumpling Run @Spring	7.8	7.6	7.5	7.5	287	254	257	273	10.3	10.6	10.5	9.4
Dumpling Run Bottom	8.1	7.8	7.6	8.3	284	264	272	291	10.4	10.6	10.3	9.7
Spring Run @Spring	7.9	7.7	7.7	8.1	296	330	358	331	10.3	10.5	10.1	9.8
Spring Run Middle	7.8	7.7	7.5	7.9	255	255	254	255	10.3	10.4	10.3	9.6
Spring Run Bottom	7.5	7.6	7.5	7.9	248	253	255	256	10.6	10.7	10.4	9.6

**Statistical comparisons:** pH was statistically lower in Spring Run Bottom than Spring Run @Spring and Dumpling Run Bottom. Conductivity was higher in Spring Run @Spring than Spring Run Bottom. No difference between sites for dissolved oxygen.


**Pre-Post Comparisons:** pH was statistically lower at Dumpling Run @Spring during the Post treatment period; Dissolved oxygen was statistically lower at all sites except Dumpling Run @Spring during the Post period.

Median values for total phosphorus (TP) and total nitrogen (TN) by site and year are provided in Table 2. Source water TP was similar in each stream, and did not increase in the downstream site in the control stream


(Dumpling Run). Both sites below the hatchery in Spring Run (Spring Run Middle and Spring Run Bottom) had significantly higher median TP than all other locations. No difference in TP between pre-treatment (2005 and 2006) and post-treatment (June 2007 through September 2008) periods was evident.

**Table 2. Median total phosphorus (TP) and total nitrogen (TN) by site and year.**

Site	TP (mg/L) Median				TN (mg/L) Median			
	2005	2006	2007	2008	2005	2006	2007	2008
Dumpling Run @Spring = DR2	0.028	0.054	0.041	0.028	0.341	0.36	0.31	0.37
Dumpling Run Bottom = DR1	0.026	0.044	0.038	0.031	0.364	0.36	0.34	0.59
Spring Run @Spring = SR3	0.025	0.049	0.036	0.038	0.641	0.57	0.52	0.69
Spring Run Middle = SR2	0.075	0.103	0.106	0.068	0.887	0.63	0.78	0.88
Spring Run Bottom = SR1	0.087	0.103	0.085	0.070	0.877	0.73	0.75	0.96



SR1 and SR2 had significantly higher total phosphorus than all other sites. There was no difference between Pre and Post treatment periods.




DR1 Pre and Post had significantly lower TN than SR1 Post. No other significant differences were detected.

Median total nitrogen was consistently but not significantly higher in the Spring Run source water than in Dumpling Run. Median TN did not increase in the downstream direction in the control stream (Dumpling


Run). Both sites below the hatchery in Spring Run (Spring Run Middle and Spring Run Bottom) had higher (but not significantly) TN than control sites in Dumpling Run and the source water in the Spring Run spring. No difference in TN between pre-treatment and post-treatment periods was evident.

**Table 3. Median BOD5 and TSS by site and year.**

Site	Median BOD5 (mg/L)				Median TSS (mg/L)			
	2005	2006	2007	2008	2005	2006	2007	2008
Dumpling Run @Spring = DR2	1.54	1.4	0.68	0.9	4.5	1.15	2.5	2
Dumpling Run Bottom = DR1	1.515	1.1	0.605	0.795	2.08	5.5	4.5	4.5
Spring Run @Spring = SR3	0.985	0.645	0.415	0.465	1.58	2.58	3.5	1.575
Spring Run Middle = SR2	0.91	0.76	0.53	0.735	5.5	5	3	2.5
Spring Run Bottom = SR1	1.01	0.425	0.415	0.54	6.5	7	6	3.5



DR1-Pre significantly higher than all but DR2-Pre and SR2-Pre.



No significant differences.



Median values for BOD5 and TSS by site and year are provided in Table 3 and graphic comparisons in the associated graphs. Source water BOD5 was distinctly (but not significantly) higher in Dumpling Run than

Spring Run. There was no marked change in BOD5 in the downstream direction in either stream. However, median BOD5 decreased from Pre to Post periods at all sites, significantly only in DR1 and DR2. TSS was similar in the source water for the two streams, with data ranging broadly. Median TSS tended to increase slightly (but not significantly) in a downstream direction in both streams.

### How did water quality vary over time?

The following four time-series bar graphs and associated text show how total N, total P, TSS and BOD5 concentrations varied during the study period to date. Also shown on each graph is the average of the flows at the two flow stations for each sampling period; this was done for the sake of graphic simplification, justified because these values were very strongly correlated ( $r^2 = 0.94$ ).

**Time series bar graphs of total nitrogen, total phosphorus concentrations at all permanent study sites.**





Total nitrogen (TN) varied widely and, based on correlation analysis, generally tracked with flows at all sites. The highest levels at all sites were observed in August 2005 during a high water event, with the exception of SpringRunBottom in April 2008, which was high due to a very high TKN (2.74 mg/l). TN was usually higher in all Spring Run sites than Dumpling Run. Elevated TN at SR Middle and SR Bottom in September 2006 was probably due to sampling that occurred on hatchery cleanout day. However,

sampling in April 2007 also occurred on a cleanout day, and a similar increase in TN at SR Middle and SR Bottom was not observed. TN remained elevated, relative to other sites, in the two point source impacted Spring Run sites after the treatment system became operational in June 2007.

Total phosphorus (TP) varied widely over time at all sites and did not apparently vary with flow levels. However, the highest TP concentrations at all sites except SR Middle and Bottom were recorded during an active runoff event in April 2006. The highest TP concentration at SR Middle occurred when sampling occurred on a hatchery cleanout day in April 2007. The highest TP concentrations at SR Bottom occurred on September 2005, May and June 2006, for no apparent reason. Elevated TP downstream of the hatchery was evident at all flows at SR Middle and Bottom (with a notable exception in August 2006 at SR Bottom). TP concentrations at the two point source sites were often distinctly different in 2006; this was not the case in 2005, 2007, and 2008. Elevated TP at SR Middle and SR Bottom in September 2006 may have been due to sampling that occurred on hatchery cleanout day; however, TP was similarly high at these two sites in September 2005 when cleanout was not occurring. TP remained elevated, relative to other sites, in the two point source impacted Spring Run sites after the treatment system became operational in June 2007.

**Time series bar graphs of total suspended solids, and biochemical oxygen demand concentrations at all permanent study sites.**




TSS varied widely and very roughly tracked with flows at all sites. The highest TSS levels for all sites were observed in August '05 during a high water event, but were not notably high during an active runoff event in April 2006. TSS concentrations were more

consistently elevated at DR Lower, SR Middle and SR Bottom in 2006 than 2005. Elevated TSS at SR Middle and SR Bottom in September 2006 and April 2007 may have been due to sampling on hatchery cleanout day; however, TSS was also high at both spring sampling sites in September 2006. TSS was similarly elevated at all sites during the low water period from July through September 2007.


BOD5 varied substantially between sites. Correlation analysis indicated that BOD5 at Spring Run point source impacted sites tended to vary with flow, while patterns of BOD5 concentrations in non point sites had no apparent relationship to flow. BOD5 concentrations were notably low during the one active runoff event in April 2006, and notably high at all sites except SR Spring during a high water event in August 2005.

**How did loads of key parameters vary over time?** The following four time-series bar graphs and associated text show how total N, total P, TSS and BOD5 loads (in pounds per hour) varied at the two flow station sites during the baseline and the June-Sept 2007 post treatment sampling period.

**Time series bar graphs of total nitrogen and total phosphorus loads (in pounds per hour) at the two flow sites.**




Total nitrogen loads varied widely and generally tracked with flows at all sites. As with TN concentrations, SR consistently had the higher TN loads. The highest loads at both sites were delivered during the three highest water events.




Total phosphorus loads varied much more widely over time in DR than SR. As with TP concentrations, SR consistently had the higher TP loads.

Time series bar graphs of total suspended solids and biochemical oxygen demand loads (in pounds per hour) at the two flow sites.



TSS loads varied widely and roughly tracked with flows at both sites. The highest loads were observed in August 2005 during a high water event. SR consistently had the higher TSS loads.




BOD5 loads varied substantially between the two sites and roughly varied with flows. The highest loads at both sites were observed in August 2005 during a high water event.

## Benthic Macroinvertebrate & Periphyton Analysis


An assessment of Spring Run in 2003 by WVDEP (Tim Craddock, 2003) collected benthic invertebrate samples at sites near those chosen for the current study. The study found low diversity at the lower station, where the most abundant family was the Chironomidae, an indicator of organic pollution. It also found abundant Gammaridae amphipods at all sites. (See Appendix 3 for results, as well as a commentary of the challenge of assessing Spring Run macroinvertebrates by WVDEP's Tim Craddock.)

Benthic samples for this project were collected twice each year, in the spring and in the autumn, at all water quality sampling sites. The methods used were the standard collection, processing and identification protocols of WV DEP, with a 200 count subsample. Summary data is provided in Appendix 4. Observations during benthic field collections indicated abundance, often overwhelming abundance, of amphipods in both streams (Craddock and Gillies, personal observations). Amphipods are often extremely abundant in limestone spring fed streams, and their abundance renders many standard benthic invertebrate indices less meaningful in assessing this type of stream. Numerical results indicate that the most abundant organisms, at all study sites, were either Gammaridae (amphipods) or Chironomidae (midges). The result of this abundance, combined with use of the standard 200 count subsampling method, often means that the actual diversity of organisms in a sample is masked.


This section focuses on four indices that were chosen for their utility in this setting: Percent EPT (Ephemeroptera - mayfly, Plecoptera - stonefly, Trichoptera - caddisfly); percent dominance; percent Gammaridae (amphipods); and percent Chironomidae (non biting midges). Discussion of results are presented in the box below the graph for each index.








Percent dominance is common metric where high numbers typically indicate poor water quality. As noted above, however, such metrics are problematic in limestone spring fed streams where dominance by amphipods (or isopods) is normal. All sites had relatively high dominance throughout the study period, with all sites having at least one incidence of >75% dominance. DR@Spring always had % dominance greater than 50.



Percent Gammaridae is not a standard metric. It is used here in recognition that amphipods are commonly abundant in limestone spring fed streams, are certainly very abundant in all the study sites, and their abundance makes a number of standard metrics unreliable. % Gammaridae varied widely at all sites, with no universal pattern apparent. A quick comparison of the % Gammaridae graph to the % Dominance graph indicates that amphipods were, in most cases, the dominant organism at every site except SR Middle. It is notable that amphipods became proportionately much more important at SR Middle from Spring 2007 through Fall 2008 than previously.



Percent Chironomidae is a common metric where high numbers typically indicate poor water quality and organic pollution. Chironomids were the dominant group at SR Middle during the first two years of this project, supplanting the amphipods that were dominant at all other sites. This site was distinctive for the large amount of organic matter and matted algae entrained in the stream sediment. Chironomids became proportionately less important at SR Middle from Spring 2007 through Fall 2008 than previously.

### Fisherman Survey

Anglers with permits to fly fish, catch-and-release were invited, by a notice posted at the Spring Run parking area, to report the date fished, species, length, and stream location of their catch. The fly-fishing, catch-and-release section of Spring Run extends for about ¾ mile. This section was arbitrarily divided into 10 sections, marked at streamside; Numbered 0 thru 9, beginning with 0 at the downstream boundary and increasing upstream. Sections were not of equal length. Anglers fished wherever they chose. Fishing sessions ranged from less than an hour to several hours. Anglers reported on a card designed with stream sections vs. 6 length categories, in inches; 0-7, 8-10, 11-13, 14-16, 17-19, 20-up. This card was available from a box located convenient to the parking area and next to a locked box for depositing completed reports. The parking area was adjacent to stream section Number 4. A member of the monitoring team collected reports frequently and summarized data monthly. The purpose of the study was to acquire data on number, size, and location of Spring Run trout, not to evaluate angler success.

Anglers cooperated in collecting data with a participation rate estimated above 80% for sessions fished. Summary data presented above are for April through December in 2005, January through December in 2006, 2007, and 2008. The most heavily fished period is April through September. Data presented in the following tables are for rainbow trout. A small number of brown, brook and golden trout were reported.

<b>Table 6. Spring Run angler catch reports.</b>												
<b>Spring Run Angler Catch Reports, Rainbow Trout: April thru Dec 2005</b>												
<b>65 Anglers Reporting      230 Fishing Sessions</b>												
<b>Length</b>	<b>Stream Section</b>										<b>Total</b>	<b>%</b>
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>		
0-7	70	108	77	130	220	335	201	142	58	47	<b>1388</b>	37.5
8--10	22	35	26	72	146	221	191	217	203	162	<b>1295</b>	35
11--13	7	5	17	27	39	75	75	89	170	175	<b>679</b>	18.3
14--16		1		16	25	23	33	27	29	86	<b>240</b>	6.5
17--19				1	5	4	9	7	10	24	<b>60</b>	1.6
20--up			1		1	4	7	10	6	13	<b>42</b>	1.1
<b>Total</b>	<b>99</b>	<b>149</b>	<b>121</b>	<b>246</b>	<b>436</b>	<b>662</b>	<b>516</b>	<b>492</b>	<b>476</b>	<b>507</b>	<b>3704</b>	
<b>%</b>	2.7	4	3.3	6.6	11.8	17.9	13.9	13.3	12.9	13.7		
<b>16.1 rainbow trout/angler session</b>												

<b>Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2006</b>												
<b>76 Anglers Reporting      232 Fishing Sessions</b>												
<b>Length</b>	<b>Stream Section</b>										<b>Total</b>	<b>%</b>
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>		
0-7	25	46	42	89	134	153	112	46	30	33	718	31.6
8--10	18	14	20	49	103	109	121	134	64	66	698	30.7
11--13	4	10	18	18	34	46	77	104	109	136	536	23.6
14--16		4	4	8	9	18	31	42	43	92	251	11
17--19		1	1	1	3	2	2	9	4	19	42	1.8
20--up				1	3	1		1	1	8	15	0.7
<b>Total</b>	<b>47</b>	<b>75</b>	<b>85</b>	<b>160</b>	<b>286</b>	<b>329</b>	<b>343</b>	<b>336</b>	<b>251</b>	<b>354</b>	<b>2272</b>	
<b>%</b>	2.1	3.3	3.7	7	12.6	14.5	15.1	14.8	11	15.6		
<b>9.8 rainbow trout/angler session</b>												


<b>Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2007</b>												
<b>59 Anglers Reporting      211 Fishing Sessions</b>												
<b>Length</b>	<b>Stream Section</b>										<b>SUMS</b>	<b>%</b>
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>		
<b>0-7</b>	20	54	30	51	61	89	51	37	27	22	<b>442</b>	<b>29.3%</b>
<b>8-10</b>	8	19	18	35	65	88	38	36	29	48	<b>384</b>	<b>25.4%</b>
<b>11-13</b>	3	9	8	15	65	62	63	53	36	76	<b>390</b>	<b>25.8%</b>
<b>14-16</b>			5	8	30	27	36	54	29	55	<b>244</b>	<b>16.2%</b>
<b>17-19</b>		1		2	7	3	2	4	2	14	<b>35</b>	<b>2.3%</b>
<b>20-up</b>						2	2	2	1	7	<b>14</b>	<b>0.9%</b>
<b>SUMS</b>	<b>31</b>	<b>83</b>	<b>61</b>	<b>111</b>	<b>228</b>	<b>271</b>	<b>192</b>	<b>186</b>	<b>124</b>	<b>222</b>	<b>1509</b>	
<b>% by Section</b>	<b>2.1%</b>	<b>5.5%</b>	<b>4.0%</b>	<b>7.4%</b>	<b>15.1%</b>	<b>18.0%</b>	<b>12.7%</b>	<b>12.3%</b>	<b>8.2%</b>	<b>14.7%</b>		
<b>7.2 rainbow trout/angler session</b>												

Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2008												
45 Anglers Reporting 171 Fishing Sessions												
Length	Stream Section										SUMS	%
	0	1	2	3	4	5	6	7	8	9		
0-7	24	64	42	62	69	118	68	87	48	72	654	47.2%
8-10	15	51	31	37	63	63	48	47	37	86	478	34.5%
11-13	1	11	8	10	22	17	7	13	33	43	165	11.9%
14-16		1	2	4	7	7	3	2	7	35	68	4.9%
17-19			1		1	1				11	14	1.0%
20-up					2			2	1	3	8	0.6%
<b>SUMS</b>	<b>40</b>	<b>127</b>	<b>84</b>	<b>113</b>	<b>164</b>	<b>206</b>	<b>126</b>	<b>151</b>	<b>126</b>	<b>250</b>	<b>1387</b>	
<b>% by Section</b>	<b>2.9%</b>	<b>9.2%</b>	<b>6.1%</b>	<b>8.1%</b>	<b>11.8%</b>	<b>14.9%</b>	<b>9.1%</b>	<b>10.9%</b>	<b>9.1%</b>	<b>18.0%</b>		

8.1 rainbow trout/angler session

In 2005, 65 anglers reported 230 fishing sessions in 2005, 76 anglers reported 232 fishing sessions in 2006, 59 anglers reported 210 fishing sessions in 2007, and 45 anglers reported 171 fishing sessions in 2008. The number of trout caught per session declined from 16.1 to 9.8 to 7.2 (in 2005, 2006, and 2007, respectively), and rose slightly to 8.1 per session in 2008.

The total catch by stream section is summarized in the following graph. In each sampling year, the greatest numbers were caught in sections “4” through “9.” The catch in every section declined sharply from a high in 2005.



## DISCUSSION

The two study streams are impacted by a variety of potential sources of pollution, some readily apparent and some not. The Spring Run watershed contains the trout rearing facility, which is a known source of BOD, TSS and nutrients, as well as a number of non point sources including poultry houses, residences, roads, and occasional cattle. The Dumpling Run watershed has no point sources, and apparently no poultry houses, but

includes residences and small farms with livestock, as well as a dirt and gravel road. In addition, the source springs in both watersheds both originate in limestone and sandstone strata and show rapid changes (turbidity, increase in flow) following heavy precipitation; this is indicative of solution channel connections through limestone at the surface of the ground.

Despite the wealth of confounding variables and within parameter variability, some patterns are reasonably clear from the above data. The spring source water for the two streams has similar pH, conductivity, dissolved oxygen, TSS, and phosphorus, and supports similar communities of benthic macroinvertebrates. Source water in Dumpling Run tends to have less nitrate, and total N than Spring Run, and higher BOD5. Conductivity and pH tend to increase or not change in a downstream direction in Dumpling Run, and tend to decrease in a downstream direction in Spring Run. Nutrients and TSS are generally similar in the two Dumpling Run sites, and tend to increase in a downstream direction in Spring Run, often dramatically.

Installation of the treatment facility did not appear to have an impact on the water quality parameters measured on non-cleanout days. Review of post-treatment water quality data indicates that the plant upgrade did not change the water quality characteristics of Spring Run's water downstream of the plant on non-cleanout days. Phosphorus in particular remains elevated. TSS and BOD5 in Spring Run remain lower, on non cleanout days, than Dumpling Run. However, it is also clear that our sampling protocol is not capturing reductions in the pollutant plume that may have occurred during cleanout as a result of the new effluent treatment system (see Hatchery Upgrade section above). Data provided by WVDNR indicates that this process reduces pollutant loads related to cleanout by roughly 90%. For example, TP concentrations in the effluent stream during cleanout fell from an average of 4.5 mg/L to 0.54 mg/L.

Benthic macroinvertebrate communities in all sites are defined by dominance of either amphipods or chironomidae midge larvae. The natural dominance of amphipods in limestone streams may well have masked underlying benthic macroinvertebrate diversity. Diversity is higher in the lower Dumpling run site than in the two point source impacted sites in Spring Run.

The change in relative abundance of chironomidae in Spring Run at the top of the managed fishing section is, perhaps, a sign that large reductions in discharge of effluent solids due to installation of the hatchery treatment system is having a positive impact on the benthic macroinvertebrate community in Spring Run. A related indicator is that the subjective condition of the stream bottom at that site was notably improved (less odor, less "greasy feel" to sediment, less entrained algae) when last sampled in the fall of 2008 (Craddock & Gillies, Personal Communication).

Decline in the Spring Run fishery was apparent throughout the study period. While there were some signs of improved Rainbow Trout spawning and recruitment in late 2008 through early 2009, it is too early to know if any improvement can be expected.

### **Erosion Control Projects**

Friends of Spring Runs Wild Trout, supported by a WV Stream Partners grant, partnered with the WVCA to reduce channel erosion and the resulting sedimentation problem in Spring Run. In 2006 WVCA supplied the design and heavy equipment to reshape and seed 210 feet of severely eroding roadside channel immediately above the spring source of Spring Run. In 2007 Friends of Spring Runs Wild Trout, with volunteered design, labor and equipment, installed three sediment basins and numerous sediment check dams and sections of rip-rap in three storm water runoff ditches, totaling about 600 feet, from developed areas. Sediment check dams of wood construction were placed in four woodland hollows which funnel run-off into Spring Run. Tree seedlings were planted in work areas in early 2008.

Landowners remain concerned about sediment entering Spring Run from public roadway ditches. It is the hope of the working group and Friends of Spring Run's Wild Trout that this will become more of a priority

for WV Division of Highways in the coming months. Cattle have been observed within the stream and along the unstable banks of the Spring Run above the fly fishing section.

### **Outreach**

A paper on the Spring Run project was presented at the 2007 Virginia/West Virginia Water Research Symposium in November 2007 by Cacapon Institute. This paper was coauthored by Neil Gillies (CI), Carla Hardy (WVCA), and Tim Craddock (WVDEP).

### **Priority Watershed Planning**

The Mill Creek Watershed, of which Spring Run is a part, was listed by the WVDEP as impaired for excess fecal coliform bacteria in 1996. Following a Total Maximum Daily Load (TMDL) study, EPA called for a 37% reduction in fecal coliform loadings. A watershed based plan to implement the TMDL is in development and will be submitted to EPA during the winter of 2008. Mill Creek is also a priority watershed for West Virginia's Potomac Tributary Strategy's Implementation Program, a Chesapeake Bay Program initiative.

A working group comprised of representatives from West Virginia Conservation Agency, West Virginia Department of Agriculture, Cacapon Institute, USDA Natural Resources Conservation Service and WVU Extension Service has worked to identify areas within the watershed that would benefit from BMP installation. A voluntary survey was mailed to all landowners within the entire watershed during the spring of 2007 requesting that they rank their environmental concerns. This information has been tallied and the working group has plans to bring educational programs that address these concerns to the local community as the project progresses.

The above steps all contribute to a process that will bring funds to the watershed for the purpose of improving water quality in the Mill Creek watershed. The funds will be used for projects such as feedlot relocations, installation of streambank fencing, alternative livestock water development, riparian buffers, failing septic upgrades, and wetland restoration. Funding for this project is anticipated in the summer of 2009.

### **Post Study Expectations**

Benthic macroinvertebrate sampling will continue for at least one year to determine if apparent improvements in Spring Run below the hatchery are sustained over time. Fisherman will continue to report "catch and release" results for the foreseeable future, and WVDNR will conduct additional electro-shocking in the future, when possible.

Appendix 1. Water Quality Statistics by Year.

Site	Yr	Minimum	Median	Maximum	Mean	Std.Dev.
<b>Ammonia-N (mg/L)</b>						
Dumpling Run @Spring	2005	0.003	0.007	0.082	0.025	0.033
Dumpling Run Bottom		0.003	0.007	0.079	0.028	0.037
Spring Run @Spring		0.003	0.012	0.915	0.167	0.367
Spring Run Middle		0.051	0.093	0.214	0.107	0.063
Spring Run Bottom		0.017	0.043	0.161	0.070	0.059
Dumpling Run @Spring	2006	0.008	0.008	0.008	0.008	0.000
Dumpling Run Bottom		0.008	0.008	0.008	0.008	0.000
Spring Run @Spring		0.008	0.008	0.008	0.008	0.000
Spring Run Middle		0.008	0.046	0.102	0.048	0.034
Spring Run Bottom		0.008	0.008	0.041	0.019	0.017
Dumpling Run @Spring	2007	0.003	0.003	0.003	0.003	0.000
Dumpling Run Bottom		0.003	0.003	0.003	0.003	0.000
Spring Run @Spring		0.003	0.003	0.026	0.007	0.009
Spring Run Middle		0.029	0.048	0.283	0.090	0.099
Spring Run Bottom		0.003	0.003	0.153	0.028	0.061
Dumpling Run @Spring	2008	0.003	0.003	0.013	0.006	
Dumpling Run Bottom		0.003	0.003	0.039	0.011	
Spring Run @Spring		0.003	0.003	0.008	0.004	
Spring Run Middle		0.003	0.066	0.214	0.081	
Spring Run Bottom		0.003	0.003	0.046	0.013	
<b>Nitrate-N (mg/L)</b>						
Dumpling Run @Spring	2005	0.17	0.23	0.38	0.26	0.079
Dumpling Run Bottom		0.19	0.26	0.50	0.31	0.117
Spring Run @Spring		0.37	0.48	0.59	0.48	0.083
Spring Run Middle		0.43	0.49	1.14	0.63	0.273
Spring Run Bottom		0.50	0.61	1.23	0.70	0.275
Dumpling Run @Spring	2006	0.12	0.16	0.36	0.19	0.088
Dumpling Run Bottom		0.13	0.18	0.40	0.21	0.097
Spring Run @Spring		0.34	0.41	0.55	0.41	0.079
Spring Run Middle		0.39	0.41	0.57	0.45	0.074
Spring Run Bottom		0.42	0.48	0.62	0.50	0.078
Dumpling Run @Spring	2007	0.00	0.16	0.22	0.14	0.077
Dumpling Run Bottom		0.02	0.18	0.25	0.16	0.082
Spring Run @Spring		0.02	0.39	0.41	0.33	0.154
Spring Run Middle		0.00	0.51	0.69	0.45	0.234
Spring Run Bottom		0.02	0.59	0.63	0.49	0.233
Dumpling Run @Spring	2008	0.12	0.15	0.26	0.17	
Dumpling Run Bottom		0.13	0.16	0.31	0.19	
Spring Run @Spring		0.36	0.42	0.58	0.44	
Spring Run Middle		0.48	0.52	0.64	0.55	
Spring Run Bottom		0.62	0.65	0.71	0.65	
<b>Nitrite-N (mg/L)</b>						
Dumpling Run @Spring	2005	0.001	0.001	0.005	0.001	0.002
Dumpling Run Bottom		0.001	0.001	0.014	0.003	0.006
Spring Run @Spring		0.001	0.001	0.003	0.001	0.001

Spring Run Middle		0.001	0.007	0.023	0.009	0.009
Spring Run Bottom		0.001	0.006	0.029	0.01	0.011
Dumpling Run @Spring	2006	0.001	0.001	0.001	0.001	0
Dumpling Run Bottom		0.001	0.001	0.001	0.001	0
Spring Run @Spring		0.001	0.001	0.001	0.001	0
Spring Run Middle		0.004	0.009	0.031	0.014	0.01
Spring Run Bottom		0.004	0.007	0.016	0.008	0.004
Dumpling Run @Spring		2007	0.001	0.001	0.001	0.001
Dumpling Run Bottom	0.001		0.001	0.001	0.001	0
Spring Run @Spring		0.001	0.001	0.001	0.001	0
Spring Run Middle		0.001	0.01	0.018	0.01	0.006
Spring Run Bottom		0.001	0.003	0.006	0.003	0.003
Dumpling Run @Spring		2008	0.001	0.001	0.001	0.001
Dumpling Run Bottom	0.001		0.001	0.001	0.001	
Spring Run @Spring		0.001	0.001	0.001	0.001	
Spring Run Middle		0.001	0.013	0.031	0.014	
Spring Run Bottom		0.001	0.001	0.018	0.007	
<b>TKN (mg/L)</b>						
Dumpling Run @Spring	2005	0.041	0.115	0.758	0.24	0.273
Dumpling Run Bottom		0.081	0.108	1.27	0.33	0.471
Spring Run @Spring		0.099	0.15	1.89	0.44	0.711
Spring Run Middle		0.214	0.305	1.46	0.51	0.475
Spring Run Bottom		0.167	0.291	0.938	0.38	0.291
Dumpling Run @Spring		2006	0.035	0.203	0.233	0.163
Dumpling Run Bottom	0.135		0.175	0.294	0.2	0.068
Spring Run @Spring		0.091	0.214	0.287	0.201	0.066
Spring Run Middle		0.181	0.214	1.09	0.387	0.355
Spring Run Bottom		0.156	0.277	0.642	0.307	0.176
Dumpling Run @Spring		2007	0.117	0.167	0.196	0.161
Dumpling Run Bottom	0.041		0.17	0.419	0.217	0.148
Spring Run @Spring		0.089	0.149	0.243	0.155	0.051
Spring Run Middle		0.167	0.288	0.527	0.302	0.122
Spring Run Bottom		0.069	0.216	0.304	0.196	0.083
Dumpling Run @Spring		2008	0.009	0.109	0.233	0.111
Dumpling Run Bottom	0.009		0.083	0.341	0.150	
Spring Run @Spring		0.009	0.131	0.271	0.145	
Spring Run Middle		0.094	0.274	1.090	0.434	
Spring Run Bottom		0.052	0.069	0.344	0.169	
<b>Total N (mg/L)</b>						
Dumpling Run @Spring	2005	0.252	0.341	1.143	0.5	0.348
Dumpling Run Bottom		0.274	0.364	1.784	0.63	0.583
Spring Run @Spring		0.476	0.641	2.453	0.93	0.755
Spring Run Middle		0.71	0.887	2.616	1.14	0.736
Spring Run Bottom		0.688	0.877	2.197	1.09	0.574
Dumpling Run @Spring		2006	0.176	0.364	0.569	0.348
Dumpling Run Bottom	0.296		0.356	0.695	0.407	0.146
Spring Run @Spring		0.522	0.57	0.838	0.615	0.116
Spring Run Middle		0.58	0.634	1.631	0.849	0.412
Spring Run Bottom		0.601	0.734	1.216	0.818	0.227



Dumpling Run @Spring	2007	0.197	0.312	0.405	0.305	0.073
Dumpling Run Bottom		0.146	0.336	0.64	0.38	0.211
Spring Run @Spring		0.165	0.517	0.644	0.485	0.167
Spring Run Middle		0.306	0.775	1.01	0.766	0.258
Spring Run Bottom		0.25	0.747	0.899	0.692	0.232
Dumpling Run @Spring	2008	0.160	0.259	0.370	0.280	
Dumpling Run Bottom		0.170	0.274	0.652	0.341	
Spring Run @Spring		0.370	0.640	0.694	0.581	
Spring Run Middle		0.735	0.837	1.631	1.002	
Spring Run Bottom		0.700	0.763	1.012	0.830	
<b>TP (mg/L)</b>						
Dumpling Run @Spring	2005	0.007	0.028	0.059	0.028	0.019
Dumpling Run Bottom		0.007	0.026	0.052	0.028	0.019
Spring Run @Spring		0.013	0.025	0.046	0.028	0.014
Spring Run Middle		0.049	0.075	0.166	0.086	0.046
Spring Run Bottom		0.059	0.087	0.14	0.092	0.031
Dumpling Run @Spring	2006	0.022	0.054	0.124	0.06	0.035
Dumpling Run Bottom		0.02	0.044	0.134	0.052	0.041
Spring Run @Spring		0.026	0.049	0.261	0.08	0.09
Spring Run Middle		0.081	0.103	0.179	0.122	0.044
Spring Run Bottom		0.012	0.103	0.143	0.094	0.053
Dumpling Run @Spring	2007	0.033	0.041	0.046	0.039	0.005
Dumpling Run Bottom		0.026	0.038	0.047	0.037	0.008
Spring Run @Spring		0.033	0.036	0.068	0.043	0.014
Spring Run Middle		0.072	0.106	0.293	0.128	0.083
Spring Run Bottom		0.068	0.085	0.101	0.084	0.012
Dumpling Run @Spring	2008	0.013	0.042	0.049	0.036	
Dumpling Run Bottom		0.039	0.042	0.049	0.042	
Spring Run @Spring		0.023	0.033	0.046	0.034	
Spring Run Middle		0.049	0.143	0.176	0.129	
Spring Run Bottom		0.029	0.124	0.140	0.104	
<b>TSS (mg/L)</b>						
Dumpling Run @Spring	2005	1.15	4.50	45.00	12.88	17.423
Dumpling Run Bottom		1.15	2.08	43.00	9.58	16.588
Spring Run @Spring		1.00	1.58	78.00	14.38	31.175
Spring Run Middle		1.15	5.50	81.00	17.36	31.281
Spring Run Bottom		4.00	6.50	72.00	18.50	26.629
Dumpling Run @Spring	2006	1.15	1.15	4.00	1.63	1.164
Dumpling Run Bottom		3.00	5.50	8.00	5.50	2.258
Spring Run @Spring		1.15	2.58	11.00	4.08	3.933
Spring Run Middle		3.00	5.00	28.00	8.67	9.522
Spring Run Bottom		3.00	7.00	19.00	8.50	5.431
Dumpling Run @Spring	2007	0.80	2.50	5.00	2.43	1.597
Dumpling Run Bottom		0.80	4.50	7.00	3.97	2.246
Spring Run @Spring		2.00	3.50	6.00	3.67	1.633
Spring Run Middle		0.80	3.00	28.00	6.80	10.442
Spring Run Bottom		3.00	6.00	7.00	5.83	1.472
Dumpling Run @Spring	2008	1.15	3.00	6.00	3.06	
Dumpling Run Bottom		1.15	3.00	6.00	3.23	

Spring Run @Spring		1.15	1.15	11.00	3.69	
Spring Run Middle		1.15	3.00	28.00	8.43	
Spring Run Bottom		2.00	6.00	7.00	5.00	
<b>Turbidity (NTU)</b>						
Dumpling Run @Spring	2005	0.45	0.9	22.95	7.52	10.6
Dumpling Run Bottom		1.24	2.12	43.8	10.29	16.83
Spring Run @Spring		1.03	1.95	18.42	5.22	6.84
Spring Run Middle		1.31	3.4	36	9.58	13.52
Spring Run Bottom		1.96	3.15	51.3	13.4	19.66
Dumpling Run @Spring	2006	0.67	0.97	2.15	1.14	0.59
Dumpling Run Bottom		2.71	3.54	4.31	3.61	0.58
Spring Run @Spring		3.14	4.78	7.47	4.86	1.66
Spring Run Middle		2.43	3.65	12.8	5.27	4
Spring Run Bottom		4.82	5.82	7.88	6.02	1.27
Dumpling Run @Spring	2007	1.2	1.58	6.82	2.49	2.18
Dumpling Run Bottom		3.32	3.82	6.89	4.45	1.33
Spring Run @Spring		2.27	4.98	7.27	4.88	1.72
Spring Run Middle		1.95	3.11	25.1	7.43	9.04
Spring Run Bottom		3.3	4.86	9.82	5.4	2.43
Dumpling Run @Spring	2008	1.07	1.28	1.94	1.39	
Dumpling Run Bottom		2.49	3.50	4.92	3.68	
Spring Run @Spring		2.27	2.59	7.47	3.54	
Spring Run Middle		1.54	4.35	12.80	5.02	
Spring Run Bottom		3.30	3.45	6.38	4.54	
<b>BOD5 (mg/L)</b>						
Dumpling Run @Spring	2005	1.01	1.54	3.13	1.81	0.88
Dumpling Run Bottom		1.18	1.52	2.68	1.63	0.55
Spring Run @Spring		0.86	0.99	1.58	1.07	0.26
Spring Run Middle		0.45	0.91	2.47	1.26	0.83
Spring Run Bottom		0.66	1.01	1.97	1.15	0.49
Dumpling Run @Spring	2006	0.76	1.4	1.97	1.35	0.46
Dumpling Run Bottom		0.76	1.1	1.59	1.12	0.34
Spring Run @Spring		0.3	0.65	1.76	0.77	0.53
Spring Run Middle		0.45	0.76	1.44	0.86	0.39
Spring Run Bottom		0.23	0.43	1.36	0.57	0.43
Dumpling Run @Spring	2007	0.15	0.68	1.06	0.61	0.33
Dumpling Run Bottom		0.23	0.61	1.21	0.61	0.37
Spring Run @Spring		0.15	0.42	0.83	0.42	0.24
Spring Run Middle		0.15	0.53	1.74	0.69	0.58
Spring Run Bottom		0.15	0.42	1.29	0.53	0.41
Dumpling Run @Spring	2008	0.68	0.90	1.29	0.94	
Dumpling Run Bottom		0.60	0.96	1.29	0.97	
Spring Run @Spring		0.30	0.48	0.96	0.62	
Spring Run Middle		0.30	0.71	1.44	0.76	
Spring Run Bottom		0.10	0.75	1.16	0.65	
<b>DO (mg/L)</b>						
Dumpling Run @Spring	2005	9.2	10.3	11.1	10.2	0.62
Dumpling Run Bottom		9.4	10.4	11.5	10.5	0.69
Spring Run @Spring		10	10.3	11.5	10.6	0.63

Spring Run Middle		10.2	10.3	11.4	10.6	0.53
Spring Run Bottom		10	10.6	11.2	10.6	0.48
Dumpling Run @Spring	2006	9.7	10.6	12.2	10.7	0.98
Dumpling Run Bottom		9.8	10.6	12.2	10.7	0.86
Spring Run @Spring		9.8	10.5	12.8	10.8	1.17
Spring Run Middle		9.8	10.4	12.6	10.7	1.09
Spring Run Bottom		9.9	10.7	12.6	10.9	1
Dumpling Run @Spring		2007	10.2	10.5	11.2	10.6
Dumpling Run Bottom	9.7		10.3	10.9	10.3	0.39
Spring Run @Spring	8.9		10.1	11.2	10.2	0.82
Spring Run Middle	9.6		10.3	10.9	10.3	0.49
Spring Run Bottom	9.9		10.4	11.1	10.5	0.45
Dumpling Run @Spring	2008		9.1	10.0	11.2	10.1
Dumpling Run Bottom		9.5	10.4	10.9	10.3	
Spring Run @Spring		9.7	10.1	10.4	10.0	
Spring Run Middle		9.5	10.0	10.3	9.9	
Spring Run Bottom		9.5	9.9	11.0	10.1	
<b>pH</b>						
Dumpling Run @Spring	2005	7.4	7.8	8	7.7	0.24
Dumpling Run Bottom		7.5	8.1	8.5	8	0.36
Spring Run @Spring		7.4	7.9	8.2	7.8	0.31
Spring Run Middle		7.3	7.8	8	7.8	0.25
Spring Run Bottom		7.2	7.5	7.7	7.5	0.23
Dumpling Run @Spring		2006	7.3	7.6	7.8	7.6
Dumpling Run Bottom	7.4		7.8	8.4	7.8	0.38
Spring Run @Spring	7.3		7.7	8.2	7.7	0.31
Spring Run Middle	7.3		7.7	8.4	7.7	0.42
Spring Run Bottom	7.3		7.6	8.1	7.6	0.31
Dumpling Run @Spring	2007		7.1	7.5	7.9	7.5
Dumpling Run Bottom		7.3	7.6	8.2	7.7	0.38
Spring Run @Spring		7.2	7.7	8.2	7.7	0.33
Spring Run Middle		6.9	7.5	8.1	7.5	0.44
Spring Run Bottom		6.9	7.5	8.2	7.5	0.44
Dumpling Run @Spring		2008	7.1	7.3	7.8	7.4
Dumpling Run Bottom	7.4		8.0	8.3	7.8	
Spring Run @Spring	7.3		7.8	8.2	7.8	
Spring Run Middle	6.9		7.7	8.1	7.6	
Spring Run Bottom	6.9		7.6	8.2	7.6	
<b>Conductivity (us/cm)</b>						
Dumpling Run @Spring	2005	45.8	286.9	372	260	112.41
Dumpling Run Bottom		48.1	283.5	352	257	106.74
Spring Run @Spring		64.6	296.1	390	269	109.38
Spring Run Middle		44.9	255	284	223	88.65
Spring Run Bottom		45.1	247.5	276	213	85.87
Dumpling Run @Spring		2006	39	254.2	372	239.6
Dumpling Run Bottom	44.9		263.9	352	241.3	103.48
Spring Run @Spring	49		330.3	391.1	263	153.97
Spring Run Middle	37.7		255.2	284	211.2	93.67
Spring Run Bottom	36.3		253.2	276	213	90.6

Dumpling Run @Spring	<b>2007</b>	234	257	345	271.7	39.83
Dumpling Run Bottom		253	271.5	340	283.2	31.72
Spring Run @Spring		271	358	375	347.5	38.3
Spring Run Middle		241	253.5	277	257.2	14.26
Spring Run Bottom		240	255	265	253.3	9.33
Dumpling Run @Spring	<b>2008</b>	234.0	247.0	356.0	266.6	
Dumpling Run Bottom		253.0	266.0	355.0	280.0	
Spring Run @Spring		288.0	378.0	390.0	363.4	
Spring Run Middle		241.0	255.0	279.0	256.0	
Spring Run Bottom		240.0	252.0	276.0	254.2	

**Appendix 2. Laboratory Methods for Water Quality Parameters.**

<b>Parameter</b>	<b>Method</b>
Ammonia Nitrogen	EPA 350.2
Nitrate	EPA 353.2
Nitrite	EPA 353.2
* Ortho Phosphate	HACH 8048
Total Phosphate	EPA 365.2
Total Kjeldahl Nitrogen	EPA 351.2
Total Suspended Solids	SM 2540D
* Turbidity	HACH 2100N
Biochemical Oxygen Demand 5	SM5210B

**Appendix 3. WV Save Our Streams Macroinvertebrate Assessment July 2003**

<b>Station 1</b> (catch-and-release)	<b>Station 2</b> (catch-and-release)	<b>Station 3</b> (above hatchery)
Ephemeroptera (mayflies)	Ephemeroptera (mayflies)	Ephemeroptera (mayflies)
<i>Baetidae</i> 73	<i>Ephemerellidae</i> 1	<i>Isonychiidae</i> 2
<i>Heptageniidae</i> 2	<i>Heptageniidae</i> 4	<i>Ephemerellidae</i> 3
Trichoptera (caddisflies)	<i>Baetidae</i> 45	<i>Baetidae</i> 30
<i>Rhyacophilidae</i> 2	Plecoptera (stoneflies)	Plecoptera (stoneflies)
<i>Hydropsychidae</i> 13	<i>Capniidae</i> 1	<i>Capniidae</i> 17
Diptera (true flies)	<i>Chloroperlidae</i> 1	<i>Perlodidae</i> 6
<i>Simuliidae</i> 8	<i>Perlodidae</i> 4	Trichoptera (caddisflies)
<i>Chironomidae</i> 67	Trichoptera (caddisflies)	<i>Rhyacophilidae</i> 3
Amphipoda (scuds)	<i>Glossosomatidae</i> 2	<i>Hydropsychidae</i> 17
<i>Gammaridae</i> 31	<i>Rhyacophilidae</i> 1	Coleoptera (beetles)
Total 196	<i>Hydropsychidae</i> 18	<i>Elmidae</i> 12
	Coleoptera (beetles)	<i>Psephenidae</i> 1
	<i>Elmidae</i> 4	Diptera (true flies)
	Diptera (true flies)	<i>Dixidae</i> 1
	<i>Simuliidae</i> 16	<i>Simuliidae</i> 12
	<i>Chironomidae</i> 37	<i>Chironomidae</i> 6
	Amphipoda (scuds)	Amphipoda (scuds)
	<i>Gammaridae</i> 125	<i>Gammaridae</i> 60
	Total 259	Total 170

**Appendix 4.** Assessing the Condition of the Macroinvertebrate Communities of Spring Run  
(Tim Craddock, Citizen's Monitoring Coordinator).

Integrity ratings for stations through study period							
Study periods	Poor	Marginal	Suboptimal	Optimal	Averages	RPD	Stations
2005 - 2006	7	9	4	0	49.5	30.63	All stations
2007 - 2008	0	5	14	1	67.4		
2005 - 2006	6	4	2	0	47.0	32.61	All Spring Run stations
2007 - 2008	0	4	8	0	65.3		
2005 - 2006	1	5	2	0	53.3	27.93	Dumpling Run stations
2007 - 2008	0	1	6	1	70.5		
2005 - 2006	4	4	0	0	43.3	34.20	Spring Run stations downstream of hatchery
2007 - 2008	0	4	4	0	61.2		

Station Codes	May-05	Oct-05	Jun-06	Oct-06
Spring Run (0.4)	SR1-505	SR1-1005	SR1-606	SR1-1006
Spring Run (1.6)	SR2-505	SR2-1005	SR2-606	SR2-1006
Spring Run (2.3)	SR3-505	SR3-1005	SR3-606	SR3-1006
Dumpling Run (1.4)	DR1-505	DR1-1005	DR1-606	DR1-1006
Dumpling Run (2.2)	DR2-505	DR2-1005	DR2-606	DR2-1006

Station Codes	May-07	Oct-07	Apr-08	Oct-08
Spring Run (0.4)	SR1-0507	SR1-1007	SR1-0408	SR1-1008
Spring Run (1.6)	SR2-0507	SR2-1007	SR2-0408	SR2-1008
Spring Run (2.3)	SR3-0507	SR3-1007	SR3-0408	SR3-1008
Dumpling Run (1.4)	DR1-0507	DR1-1007	DR1-0408	DR1-1008
Dumpling Run (2.2)	DR2-0507	DR2-1007	DR2-0408	DR2-1008

Codes	Total Taxa	EPT Taxa	Biotic Index	% EPT	% Dominant	% Tolerant	Stream Index	Integrity
SR1-0505	11	5	5.39	8.9	65.5	19.1	48.7	Marginal
SR2-0505	14	7	5.76	38.5	42.3	54.0	58.8	Marginal
SR3-0505	16	10	4.78	40.4	41.3	16.1	70.3	Suboptimal
DR1-0505	16	9	4.80	31.0	43.8	16.7	68.2	Suboptimal
DR2-0505	10	5	4.78	14.7	80.4	3.1	50.1	Marginal
SR1-1005	6	3	5.01	1.6	77.6	4.7	41.7	Poor
SR2-1005	11	5	7.20	4.6	76.9	78.5	31.1	Poor
SR3-1005	8	4	4.89	6.4	87.2	2.7	44.1	Poor

Codes	Total Taxa	EPT Taxa	Biotic Index	% EPT	% Dominant	% Tolerant	Stream Index	Integrity
DR1-1005	14	9	4.54	25.3	68.8	0.8	63.5	Suboptimal
DR2-1005	11	7	4.85	7.1	87.9	2.5	50.4	Marginal
SR1-0606	9	6	5.03	13.7	53.1	15.0	53.4	Marginal
SR2-0606	12	5	6.55	23.9	64.3	64.8	42.0	Poor
SR3-0606	12	8	4.62	32.8	55.9	7.8	63.5	Suboptimal
DR1-0606	7	2	5.00	1.0	87.6	1.9	39.5	Poor
DR2-0606	10	7	4.82	10.4	82.2	4.3	51.2	Marginal
SR1-1006	9	4	5.32	7.0	68.1	14.6	45.9	Marginal
SR2-1006	10	2	7.31	3.3	80.5	80.9	24.8	Poor
SR3-1006	7	2	4.90	2.2	90.2	0.4	39.4	Poor
DR1-1006	12	6	4.78	10.2	83.6	1.8	51.6	Marginal
DR2-1006	11	7	4.85	10.2	83.4	4.9	51.5	Marginal
SR1-0507	11	6	4.62	38.8	45.3	9.3	62.8	Suboptimal
SR2-0507	9	5	5.04	45.1	34.4	20.0	60.7	Suboptimal
SR3-0507	14	9	4.04	71.0	34.4	9.0	78.6	Suboptimal
DR1-0507	14	10	3.88	53.2	40.3	3.9	76.6	Suboptimal
DR2-0507	12	7	4.49	44.9	51.3	1.7	66.3	Suboptimal
SR1-1007	13	7	4.66	63.0	34.2	9.6	73.2	Suboptimal
SR2-1007	11	4	5.92	25.5	34.8	35.5	53.3	Marginal
SR3-1007	16	10	4.23	47.4	43.2	0.9	76.8	Suboptimal
DR1-1007	21	12	4.04	59.4	8.7	27.0	88.2	Optimal
DR2-1007	16	10	4.63	29.8	50.8	8.7	69.7	Suboptimal
SR1-0408	13	8	5.07	17.9	66.9	10.7	57.3	Marginal
SR2-0408	11	6	5.79	23.6	38.2	38.2	53.8	Marginal
SR3-0408	11	10	4.72	28.6	59.1	9.5	66.7	Suboptimal
DR1-0408	14	9	4.36	31.3	59.3	5.1	66.0	Suboptimal
DR2-0408	13	8	4.83	25.2	49.5	18.8	61.4	Suboptimal
SR1-1008	13	8	4.74	38.2	27.3	13.2	69.6	Suboptimal
SR2-1008	8	4	5.26	40.1	28.8	18.5	58.6	Marginal
SR3-	16	10	4.64	36.4	43.6	7.6	71.9	Suboptimal



Codes	Total Taxa	EPT Taxa	Biotic Index	% EPT	% Dominant	% Tolerant	Stream Index	Integrity
1008								
DR1-1008	15	9	3.88	51.4	37.5	2.4	76.6	Suboptimal
DR2-1008	13	9	4.52	16.6	74.6	1.5	59.5	Marginal